

2 Week Recheck – Pelvic Fracture Repair

Patient: _____

Date: _____

Incision

- Not applicable
- The skin incision has healed and no additional treatment is needed.
- The incision is irritated because your pet has been licking the site.
- The following is recommended:
 - Apply Neosporin Plus Pain control two to three times daily to the incision or irritated skin for 7 to 10 days.
 - Apply Bitter Apple solution (available at pet stores) around the incision twice daily for 7 days.
 - Keep an Elizabethan collar on your pet until the incision has healed (inflatable collars may be available at Pet Smart or your primary care veterinarian).

Exercise

- A quick-lift sling should be used when going up or down stairs or if your pet needs assistance to walk.
- A leash must be kept on your companion whenever going outdoors.
- Swimming is encouraged (10 minutes, 2 to 4 times daily).
- Ten minute slow walks are permitted 3 times daily until the next recheck (in 6 weeks from now).
- Jumping, running, unattended stair climbing and playing with other pets are not permitted.

Rehabilitation

- Range of motion exercises should be performed twice daily for 6 weeks. This involves fully extending, flexing, and abducting the hip. Hold the hip in maximal flexion, extension and abduction for 3 seconds in each position. Also do range of motion exercises of the knee and ankle. Twenty repetitions should be done during each session.
- We recommend rehabilitation therapy by a professional therapist.

Medication and Diet

- NSAID (_____ mg tablets): give _____ tab every _____ hours for _____ weeks.
- Tramadol 50 mg: give _____ tablet(s) every _____ hours for _____ weeks.
- Dasuquin or Cosequin, as directed on the bottle.

Summary

- The healing of the pelvis is progressing normally.
- The healing of the pelvis is progressing slower than normal.

Next recheck evaluation

Please make an appointment to have another evaluation performed in 6 weeks from now.